

Supplemental Information Text:

Analyzing and Quantifying the Gain-of-Function Enhancement of IP₃ Receptor Gating by Familial Alzheimer's Disease-Causing mutants in Presenilins

Don-On Daniel Mak¹, King-Ho Cheung², Patrick Toglia³, J. Kevin Foskett^{1,4},
and Ghanim Ullah^{3,*}

¹Department of Physiology, University of Pennsylvania,
Philadelphia, PA 19104, USA

²Department of Physiology, The University of Hong Kong,
Pok Fu Lam, Hong Kong

³Department of Physics, University of South Florida,
Tampa, FL 33620, USA,

⁴Department of Cell and Developmental Biology, University of Pennsylvania,
Philadelphia, PA 19104, USA

1 Stochastic Scheme of Channel Gating

The gating of IP₃R is given by the 12-state model described in the main text. To determine the state of the channel, we have to determine the transition probabilities at a given time [1, 2]. That is, if the j^{th} channel is in state i , we have to determine the probabilities with which it remains in that state or switches into another state allowed by the kinetic scheme shown in Fig. 1 (main text) within the time interval Δt . For example, if a channel is in state C_{00}^L , possible transitions are to states C_{20}^L and C_{04}^I . For a sufficiently small time interval Δt , the probabilities for these transitions are given by $P_{C_{00}^L \rightarrow C_{20}^L}^{(j)} = r_{(C_{00}^L \rightarrow C_{20}^L)} \Delta t$ and $P_{C_{00}^L \rightarrow C_{04}^I}^{(j)} = r_{(C_{00}^L \rightarrow C_{04}^I)} \Delta t$. The probability for the channel to remain in state C_{00}^L is $P_{C_{00}^L \rightarrow C_{00}^L} = 1 - P_{C_{00}^L \rightarrow C_{20}^L} - P_{C_{00}^L \rightarrow C_{04}^I}$. To determine the transition probabilities, we divide the unit interval into three subintervals of length $P_{C_{00}^L \rightarrow i} \Delta t$, i represent the three states to which the channel can make transition. If a random number drawn from a uniform distribution over the unit interval falls into the subinterval $P_{C_{00}^L \rightarrow i} \Delta t$, the corresponding transition is

performed. The time interval Δt was kept small enough for the linear dependence of $P_{i \rightarrow i}$ on the time interval to remain valid. We used a time step of $1 \mu s$ for the puff simulations. The channel is open when in any of the states O_{14}^I , O_{24}^I , or O_{24}^H . The above procedure was repeated for all channels.

2 Elements of Tridiagonal Matrix (TM)

Considering a spherical symmetry around the channel, the system of eqs. (26, main text) and (27, main text) converts to a TM system. Here we derive the elements of the TM. The detail derivation of the scheme is given in [3]. This method converts a 3D problem into a 1D problem and is numerically significantly faster than the standard methods such as Crank-Nicolson. In what follows n is the time index and j is the space index in spherical polar coordinates. Using eq. (29, main text), we can write eq. (26, main text) as

$$\frac{c^{(n+1)} - c^{(n)}}{\Delta t} = D \nabla^2 c^{(n+1)} + J \delta(r) + k_d^r (B_d - b_d^{(n)}) - k_d^f c^{(n+1)} b_d^{(n)} \quad (1S)$$

Writing the Laplacian in spherical polar coordinates and considering no-flux boundary conditions we get the elements of lower, middle, and upper diagonal (a_j , b_j , and c_j respectively) of the TM given as

$$a_j = \begin{cases} 0 & \text{if } j = 1 \\ \frac{-D\Delta t}{\Delta r^2} \left(1 - \frac{1}{2j}\right)^2 & \text{if } j = 2, \dots, N-1 \\ \frac{-2D\Delta t}{\Delta t} & \text{if } j = N. \end{cases} \quad (2S)$$

$$b_j = \begin{cases} \frac{6D\Delta t}{\Delta r^2} + 1 + k_d^f b_d^{(n,j)} \Delta t & \text{if } j = 1 \\ \frac{D\Delta t}{\Delta r^2} \left(1 + \frac{1}{2j}\right)^2 + \frac{D\Delta t}{\Delta r^2} \left(1 - \frac{1}{2j}\right)^2 + 1 + k_d^f b_d^{(n,j)} \Delta t & \text{if } j = 2, \dots, N-1 \\ \frac{2D\Delta t}{\Delta r^2} + 1 + k_d^f b_d^{(n,j)} \Delta t & \text{if } j = N. \end{cases} \quad (3S)$$

$$c_j = \begin{cases} \frac{-6D\Delta t}{\Delta r^2} & \text{if } j = 1 \\ \frac{-D\Delta t}{\Delta r^2} \left(1 + \frac{1}{2j}\right)^2 & \text{if } j = 2, \dots, N-1 \\ 0 & \text{if } j = N. \end{cases} \quad (4S)$$

The right hand side for the TM system for the cytosolic Ca^{2+} concentration is represented by d_j and is

$$d_j = c^{(n,j)} + J\delta(r)\Delta t + k_d^r(B_d - b_d^{(n,j)})\Delta t, \text{ j}=1,..N \quad (5S)$$

The rate equation for dye buffer can be expanded and solved iteratively in similar fashion.

References

1. Ullah G, Jung P (2006) Modeling the statistics of elementary calcium release events. *Biophysical Journal* 90: 3485–3495.
2. Swaminathan D, Ullah G, Jung P (2009) A simple sequential-binding model for calcium puffs. *Chaos* 19: 037109.
3. Ullah G, Parker I, Mak DOD, Pearson JE (2012) Multi-scale data-driven modeling and observation of calcium puffs. *Cell calcium* 52: 152–160.
4. Falcke M (2004) Reading the patterns in living cells: the physics of Ca^{2+} signaling. *Advances in physics* 53: 255–440.
5. Smith IF, Parker I (2009) Imaging the quantal substructure of single IP₃R channel activity during Ca^{2+} puffs in intact mammalian cells. *Proceedings of the National Academy of Sciences* 106: 6404–6409.
6. Williams A, West D, Sitsapesan R (2001) Light at the end of the Ca^{2+} -release channel tunnel: structures and mechanisms involved in ion translocation in ryanodine receptor channels. *Quarterly Reviews of Biophysics* 34: 61–104.
7. Taylor C, da Fonseca P, Morris E (2004) IP₃ receptors: the search for structure. *Trends in biochemical sciences* 29: 210–219.
8. Rüdiger S, Shuai JW, Sokolov IM (2010) Law of mass action, detailed balance, and the modeling of calcium puffs. *Phys Rev Lett* 105: 048103.
9. Allbritton N, Meyer T, Stryer L (1990) Range of messenger action of calcium ion and inositol 1, 4, 5-trisphosphate. *Genes Dev* 4: 1753.
10. Dargan S, Parker I (2003) Buffer kinetics shape the spatiotemporal patterns of IP₃-evoked Ca^{2+} signals. *The Journal of Physiology* 553: 775.
11. Shuai J, Parker I (2005) Optical single-channel recording by imaging Ca^{2+} flux through individual ion channels: theoretical considerations and limits to resolution. *Cell Calcium* 37: 283–299.

Table 1S: Parameters for puff simulations

Quantity	Symbol	—	Numerical Value	Reference
Resting Cytosolic Calcium	Ca_{rest}	=	$50nM$	[4]
Dye Buffer	B_d	=	$20\ \mu M$	
Number of channels	N_{ch}	=	10	[5]
Pore Radius	r_{pore}	=	$2.5nm$	[6, 7]
Channel Spacing	r_{nn}	=	$120nm$	[3, 8]
Ca^{2+}	D_c	=	$0.223\mu m^2/ms$	[9]
Dye	D_d	=	$0.200\mu m^2/ms$	[10]
Dye Buffer	k_d^f	=	$0.15/\mu Mms$	[10, 11]
	k_d^r	=	$0.45/ms$	[10, 11]